

UPSC Civil Services Main 1988 - Mathematics

Linear Algebra

Brij Bhooshan

Asst. Professor

B.S.A. College of Engg & Technology

Mathura

Question 1(a) Show that a linear transformation of a vector space \mathcal{V}_m of dimension m into a vector space \mathcal{V}_n of dimension n over the same field can be represented as a matrix. If \mathbf{T} is a linear transformation of \mathcal{V}_2 into \mathcal{V}_4 such that $\mathbf{T}(3, 1) = (4, 1, 2, 1)$ and $\mathbf{T}(-1, 2) = (3, 0, -2, 1)$, then find the matrix of \mathbf{T} .

Solution. Let $\mathbf{v}_i, i = 1, \dots, m$ be a basis of \mathcal{V}_m and $\mathbf{w}_j, j = 1, \dots, n$ be a basis of \mathcal{V}_n . If

$$\mathbf{T}(\mathbf{v}_i) = \sum_{j=1}^n a_{ji} \mathbf{w}_j, \quad i = 1, \dots, m$$

then \mathbf{T} corresponds to the $n \times m$ matrix \mathbf{A} whose (i, j) 'th entry is a_{ij} . In fact $(\mathbf{v}_1, \dots, \mathbf{v}_m) = (\mathbf{w}_1, \dots, \mathbf{w}_n)\mathbf{A}$.

It can be easily seen that

$$\begin{aligned} \mathbf{e}_1 &= (1, 0) = \frac{2}{7}(3, 1) - \frac{1}{7}(-1, 2) \\ \mathbf{e}_2 &= (0, 1) = \frac{1}{7}(3, 1) + \frac{3}{7}(-1, 2) \end{aligned}$$

and therefore

$$\begin{aligned}
\mathbf{T}(\mathbf{e}_1) &= \frac{2}{7}(4, 1, 2, 1) - \frac{1}{7}(3, 0, -2, 1) \\
&= \frac{1}{7}(5, 2, 6, 1) \\
&= \frac{1}{7}(5\mathbf{e}_1^* + 2\mathbf{e}_2^* + 6\mathbf{e}_3^* + \mathbf{e}_4^*) \\
\mathbf{T}(\mathbf{e}_2) &= \frac{1}{7}(4, 1, 2, 1) + \frac{3}{7}(3, 0, -2, 1) \\
&= \frac{1}{7}(13, 1, -4, 4) \\
&= \frac{1}{7}(13\mathbf{e}_1^* + \mathbf{e}_2^* - 4\mathbf{e}_3^* + 4\mathbf{e}_4^*)
\end{aligned}$$

Thus \mathbf{T} corresponds to the matrix $\frac{1}{7} \begin{pmatrix} 5 & 13 \\ 2 & 1 \\ 6 & -4 \\ 7 & 4 \end{pmatrix}$ w.r.t. the standard basis. ■

Question 1(b) If \mathcal{M}, \mathcal{N} are finite dimensional subspaces of \mathcal{V} , then show that $\dim(\mathcal{M} + \mathcal{N}) = \dim \mathcal{M} + \dim \mathcal{N} - \dim(\mathcal{M} \cap \mathcal{N})$.

Solution. Let $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r\}$ be a basis of $\mathcal{M} \cap \mathcal{N}$ where $\dim(\mathcal{M} \cap \mathcal{N}) = r$. Complete $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r\}$ to a basis $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_m\}$ of \mathcal{M} , where $\dim \mathcal{M} = m + r$. Complete $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r\}$ to a basis $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r, \mathbf{w}_1, \dots, \mathbf{w}_n\}$ of \mathcal{N} , where $\dim \mathcal{N} = n + r$. We shall show that $\mathcal{B} = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_m, \mathbf{w}_1, \dots, \mathbf{w}_n\}$ is a basis of $\mathcal{M} + \mathcal{N}$, proving the result.

If $\mathbf{u} \in \mathcal{M} + \mathcal{N}$, then $\mathbf{u} = \mathbf{v} + \mathbf{w}$ for some $\mathbf{v} \in \mathcal{M}, \mathbf{w} \in \mathcal{N}$. Since \mathcal{B} is a superset of the bases of \mathcal{M}, \mathcal{N} , \mathbf{v}, \mathbf{w} can be written as linear combination of elements of $\mathcal{B} \Rightarrow \mathbf{u}$ can be written as a linear combination of elements of \mathcal{B} . Thus \mathcal{B} generates $\mathcal{M} + \mathcal{N}$.

We now show that the set \mathcal{B} is linearly independent. If possible let

$$\sum_{i=1}^n \alpha_i \mathbf{v}_i + \sum_{i=1}^m \beta_i \mathbf{w}_i + \sum_{i=1}^r \gamma_i \mathbf{u}_i = \mathbf{0}$$

Since $\sum_{i=1}^n \alpha_i \mathbf{v}_i = -\sum_{i=1}^m \beta_i \mathbf{w}_i - \sum_{i=1}^r \gamma_i \mathbf{u}_i$ it follows that $\sum_{i=1}^n \alpha_i \mathbf{v}_i \in \mathcal{N}$. Therefore $\sum_{i=1}^n \alpha_i \mathbf{v}_i \in \mathcal{M} \cap \mathcal{N} \Rightarrow \sum_{i=1}^n \alpha_i \mathbf{v}_i = \sum_{i=1}^r \eta_i \mathbf{u}_i$ for $\eta_i \in \mathbb{R}$. This means that $\sum_{i=1}^n \alpha_i \mathbf{v}_i - \sum_{i=1}^r \eta_i \mathbf{u}_i = \mathbf{0}$. But $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_m\}$ are linearly independent, so $\alpha_i = 0, 1 \leq i \leq n$. Similarly we can show that $\beta_i = 0, 1 \leq i \leq m$. Then the linear independence of $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r\}$ shows that $\gamma_i = 0, 1 \leq i \leq r$. Thus the vectors in \mathcal{B} are linearly independent and form a basis of $\mathcal{M} + \mathcal{N}$, showing that the dimension of $\mathcal{M} + \mathcal{N}$ is $m + n + r = (m + r) + (n + r) - r$, which completes the proof. ■

Question 1(c) Determine a basis of the subspace spanned by the vectors $\mathbf{v}_1 = (1, 2, 3)$, $\mathbf{v}_2 = (2, 1, -1)$, $\mathbf{v}_3 = (1, -1, -4)$, $\mathbf{v}_4 = (4, 2, -2)$.

Solution. $\mathbf{v}_1, \mathbf{v}_2$ are linearly independent because if $\alpha\mathbf{v}_1 + \beta\mathbf{v}_2 = \mathbf{0}$ then $\alpha + 2\beta = 0, 2\alpha + \beta = 0, 3\alpha - \beta = 0 \Rightarrow \alpha = \beta = 0$. If $\mathbf{v}_3 = \alpha\mathbf{v}_1 + \beta\mathbf{v}_2$, then the three linear equations $\alpha + 2\beta = 1, 2\alpha + \beta = -1, 3\alpha - \beta = -4$ should be consistent — clearly $\alpha = -1, \beta = 1$ satisfy all three, showing $\mathbf{v}_3 = \mathbf{v}_2 - \mathbf{v}_1$. Again suppose $\mathbf{v}_4 = \alpha\mathbf{v}_1 + \beta\mathbf{v}_2$, then the three linear equations $\alpha + 2\beta = 4, 2\alpha + \beta = 2, 3\alpha - \beta = -2$ should be consistent — clearly $\alpha = 0, \beta = 2$ satisfy all three, showing $\mathbf{v}_4 = 2\mathbf{v}_2$.

Hence $\{\mathbf{v}_1, \mathbf{v}_2\}$ is a basis for the vector space generated by $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$. ■

Question 2(a) Show that it is impossible for $\mathbf{S} = \begin{pmatrix} a_1 & b \\ b & a_2 \end{pmatrix}, b \neq 0$ to have identical eigenvalues.

Solution. We know given \mathbf{S} symmetric $\exists \mathbf{O}$ orthogonal so that $\mathbf{O}'\mathbf{S}\mathbf{O} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$, where λ_1, λ_2 are eigenvalues of \mathbf{S} . If $\lambda_1 = \lambda_2$, then we have $\mathbf{S} = \mathbf{O}'^{-1}(\lambda\mathbf{I})\mathbf{O}^{-1} = \lambda(\mathbf{O}\mathbf{O}')^{-1} = \lambda\mathbf{I} \Rightarrow \mathbf{S} = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$. Thus if $b \neq 0$, \mathbf{S} cannot have identical eigenvalues. ■

Question 2(b) Prove that the eigenvalues of a Hermitian matrix are all real and the eigenvalues of a skew-Hermitian matrix are either zero or pure imaginary.

Solution. See question 2(a), year 1998. ■

Question 2(c) If $\mathbf{x}'\mathbf{A}\mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$, \mathbf{A} symmetric, then for all $\mathbf{y} \neq \mathbf{0}$ $\mathbf{y}'\mathbf{A}^{-1}\mathbf{y} > 0$. If λ is the largest eigenvalue of \mathbf{A} , then

$$\lambda = \sup_{\substack{\mathbf{x} \in \mathbb{R}^n \\ \mathbf{x} \neq \mathbf{0}}} \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{x}}$$

Solution. Clearly $\mathbf{A} = \mathbf{A}'\mathbf{A}^{-1}\mathbf{A} \therefore \mathbf{x}'\mathbf{A}\mathbf{x} = \mathbf{x}'\mathbf{A}'\mathbf{A}^{-1}\mathbf{A}\mathbf{x} = \mathbf{y}'\mathbf{A}^{-1}\mathbf{y}$ where $\mathbf{y} = \mathbf{A}\mathbf{x}$ for any $\mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq \mathbf{0}$. Since $|\mathbf{A}| \neq 0$, any vector \mathbf{y} can be written as $\mathbf{A}\mathbf{x}$, by taking $\mathbf{x} = \mathbf{A}^{-1}\mathbf{y}$. Thus $\mathbf{x}'\mathbf{A}\mathbf{x} > 0 \Rightarrow \mathbf{y}'\mathbf{A}^{-1}\mathbf{y} > 0$ for all $\mathbf{y} \neq \mathbf{0}$.

Let $M = \sup_{\substack{\mathbf{x} \in \mathbb{R}^n \\ \mathbf{x} \neq \mathbf{0}}} \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{x}}$. Let \mathbf{O} be an orthogonal matrix such that $\mathbf{O}'\mathbf{A}\mathbf{O} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$.

Let $\mathbf{0} \neq \mathbf{x} = \mathbf{O}\mathbf{y}$, then $\mathbf{x}'\mathbf{x} = \mathbf{y}'\mathbf{O}'\mathbf{O}\mathbf{y} = \mathbf{y}'\mathbf{y}$. Now $\mathbf{x}'\mathbf{A}\mathbf{x} = \mathbf{y}'\mathbf{O}'\mathbf{A}\mathbf{O}\mathbf{y} = \sum_i \lambda_i y_i^2 \leq \lambda \mathbf{y}'\mathbf{y}$ where λ is the largest eigenvalue of \mathbf{A} . Thus $\lambda \geq \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{y}'\mathbf{y}} = \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{x}}$, so $\lambda \geq M$. On the other hand, if $\mathbf{x} \neq \mathbf{0}$ is an eigenvector corresponding to λ , then $\mathbf{x}'\mathbf{A}\mathbf{x} = \lambda \mathbf{x}'\mathbf{x} \Rightarrow \lambda = \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{x}} \leq M$. Thus $\lambda = M$ as required. ■

Question 3(a) By converting \mathbf{A} to an echelon matrix, determine its rank, where

$$\mathbf{A} = \begin{pmatrix} 0 & 0 & 1 & 2 & 8 & 9 \\ 0 & 0 & 4 & 6 & 5 & 3 \\ 0 & 2 & 3 & 1 & 4 & 7 \\ 0 & 3 & 0 & 9 & 3 & 7 \\ 0 & 0 & 5 & 7 & 3 & 1 \end{pmatrix}$$

Solution. Consider

$$\mathbf{A}' = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 3 & 0 \\ 1 & 4 & 3 & 0 & 5 \\ 2 & 6 & 1 & 9 & 7 \\ 8 & 5 & 4 & 3 & 3 \\ 9 & 3 & 7 & 7 & 1 \end{pmatrix}$$

Interchange the first row with the third, then third with fourth, fourth with fifth and fifth with sixth to get

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 0 & 2 & 3 & 0 \\ 2 & 6 & 1 & 9 & 7 \\ 8 & 5 & 4 & 3 & 3 \\ 9 & 3 & 7 & 7 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Now perform $\mathbf{R}_3 - 2\mathbf{R}_1$, $\mathbf{R}_4 - 8\mathbf{R}_1$, $\mathbf{R}_5 - 9\mathbf{R}_1$ to get

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 0 & 2 & 3 & 0 \\ 0 & -2 & -5 & 9 & -3 \\ 0 & -27 & -20 & 3 & -37 \\ 0 & -33 & -20 & 7 & -44 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Interchange the second and the third row, and perform $-\frac{1}{2}\mathbf{R}_2$, $\frac{1}{2}\mathbf{R}_3$ to get

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & -27 & -20 & 3 & -37 \\ 0 & -33 & -20 & 7 & -44 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Perform $\mathbf{R}_4 + 27\mathbf{R}_2, \mathbf{R}_5 + 33\mathbf{R}_2$ to get

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & 0 & \frac{95}{2} & -\frac{237}{2} & \frac{7}{2} \\ 0 & 0 & \frac{125}{2} & -\frac{283}{2} & \frac{11}{2} \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Operation $\mathbf{R}_4 - \frac{95}{2}\mathbf{R}_3, \mathbf{R}_5 - \frac{125}{2}\mathbf{R}_3$ yields

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & 0 & 0 & -\frac{759}{2} & \frac{7}{2} \\ 0 & 0 & 0 & -\frac{941}{4} & \frac{11}{2} \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Now multiply \mathbf{R}_4 with $-\frac{4}{759}$

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & 0 & 0 & 1 & -\frac{14}{759} \\ 0 & 0 & 0 & -\frac{941}{4} & \frac{11}{2} \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Performing $\mathbf{R}_5 + \frac{941}{4}\mathbf{R}_4$ results in

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & 0 & 0 & 1 & -\frac{14}{759} \\ 0 & 0 & 0 & 0 & \frac{11}{2} - \frac{941 \times 7}{1882} \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

which can be converted to

$$\mathbf{A}' \sim \begin{pmatrix} 1 & 4 & 3 & 0 & 5 \\ 0 & 1 & \frac{5}{2} & -\frac{9}{2} & \frac{3}{2} \\ 0 & 0 & 1 & \frac{3}{2} & 0 \\ 0 & 0 & 0 & 1 & -\frac{14}{759} \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

which is an echelon matrix. Its rank is clearly 5, so the rank of $\mathbf{A} = 5$. ■

Question 3(b) Given $\mathbf{AB} = \mathbf{AC}$ does it follow that $\mathbf{B} = \mathbf{C}$? Can you provide a counterexample?

Solution. It does not follow that $\mathbf{B} = \mathbf{C}$.

$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \Rightarrow \mathbf{AB} = \mathbf{0}$$

$\mathbf{C} = \mathbf{0} \Rightarrow \mathbf{AC} = \mathbf{0}$, but $\mathbf{B} \neq \mathbf{C}$. ■

Question 3(c) Find a nonsingular matrix which diagonalizes $\mathbf{A} = \begin{pmatrix} 0 & -1 & 0 \\ -1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix}$, $\mathbf{B} =$

$\begin{pmatrix} 2 & 1 & -2 \\ 1 & 2 & -2 \\ -2 & -2 & 3 \end{pmatrix}$ simultaneously. Find the diagonal form of \mathbf{A} .

Solution.

$$|\mathbf{A} - \lambda\mathbf{B}| = 0 \Rightarrow \begin{vmatrix} -2\lambda & -1 - \lambda & 2\lambda \\ -1 - \lambda & -1 - 2\lambda & 1 + 2\lambda \\ 2\lambda & 1 + 2\lambda & -3\lambda \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} -2\lambda & -1 - \lambda & 0 \\ -1 + \lambda & 0 & 0 \\ 2\lambda & 1 + 2\lambda & -\lambda \end{vmatrix} = 0$$

Thus $\lambda = 0, 1, -1$. This shows that the matrices are diagonalizable simultaneously.

We now determine $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ such that $(\mathbf{A} - \lambda\mathbf{B})\mathbf{x}_i = \mathbf{0}, i = 1, 2, 3$. For $\lambda = 0$, let $\mathbf{x}_1' = (x_1, x_2, x_3)$ be such that $(\mathbf{A} - \lambda\mathbf{B})\mathbf{x}_1 = \mathbf{0}$. Thus

$$\begin{pmatrix} 0 & -1 & 0 \\ -1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \mathbf{0}$$

Thus $-x_2 = 0, -x_1 - x_2 + x_3 = 0, x_2 = 0$. Thus $\mathbf{x}_1' = (1, 0, 1)$.

For $\lambda = 1$, let $\mathbf{x}_2' = (x_1, x_2, x_3)$ be such that $(\mathbf{A} - \lambda\mathbf{B})\mathbf{x}_2 = \mathbf{0}$. Thus

$$\begin{pmatrix} -2 & -2 & 2 \\ -2 & -3 & 3 \\ 2 & 3 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \mathbf{0}$$

Thus $-2x_1 - 2x_2 + 2x_3 = 0, -2x_1 - 3x_2 + 3x_3 = 0, 2x_1 + 3x_2 - 3x_3 = 0 \Rightarrow x_2 - x_3 = 0 \Rightarrow x_1 = 0$.

Thus we may take $\mathbf{x}_2' = (0, 1, 1)$.

For $\lambda = -1$, let $\mathbf{x}_3' = (x_1, x_2, x_3)$ be such that $(\mathbf{A} - \lambda\mathbf{B})\mathbf{x}_3 = \mathbf{0}$. Thus

$$\begin{pmatrix} 2 & 0 & -2 \\ 0 & 1 & -1 \\ -2 & -1 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \mathbf{0}$$

Thus $2x_1 - 2x_3 = 0, x_2 - x_3 = 0, -2x_1 - x_2 + 3x_3 = 0 \Rightarrow x_1 = x_2 = x_3$. Thus we may take $\mathbf{x}_3' = (1, 1, 1)$.

Let $\mathbf{P} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ so that

$$\mathbf{P}'\mathbf{A}\mathbf{P} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 & 0 \\ -1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 & -1 \\ 0 & 0 & -1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

■