

1.5.5-2009

Sl. No.

8206

B-JGT-J-NBA

MATHEMATICS

Paper I

Time Allowed : Three Hours

Maximum Marks : 200

INSTRUCTIONS

Candidates should attempt questions 1 and 5 which are compulsory, and any THREE of the remaining questions selecting at least ONE question from each Section.

All questions carry equal marks.

Answers must be written in ENGLISH only.

Assume suitable data, if considered necessary, and indicate the same clearly.

Unless indicated otherwise, symbols & notations carry their usual meaning.

(Contd.)

Section - A

1. Answer any *four* of the following questions :

(a) Let V be the vector space of polynomials over R . Let U and W be the subspaces generated by

$$\{t^3 + 4t^2 - t + 3, t^3 + 5t^2 + 5, 3t^3 + 10t^2 - 5t + 5\}$$

and

$$\{t^3 + 4t^2 + 6, t^3 + 2t^2 - t + 5, 2t^3 + 2t^2 - 3t + 9\}$$

respectively. Find

(i) $\dim (U+W)$

(ii) $\dim (U \cap W)$. 10

(b) Find a linear map $T: R^3 \rightarrow R^4$ whose image is generated by $(1, 2, 0, -4)$ and $(2, 0, -1, -3)$.

10

(c) (i) Find the difference between the maximum and the minimum of the function

$$\left(a - \frac{1}{a} - x\right)(4 - 3x^2) \text{ where } a \text{ is a constant and greater than zero.}$$

(ii) If $f(h) = f(0) + hf'(0) + \frac{h^2}{2!} f''(\theta h)$,

$$0 < \theta < 1$$

Find θ , when $h = 1$ and $f(x) = (1-x)^{5/2}$.

$$5+5=10$$

(d) Evaluate :

$$(i) \int_0^{\pi/2} \frac{\sin^2 x \, dx}{\sin x + \cos x} \quad (ii) \int_1^{\infty} \frac{x^2 \, dx}{(1+x^2)^2} \quad 6+4=10$$

(e) Show that the plane $x + 2y - z = 4$ cuts the sphere $x^2 + y^2 + z^2 - x + z = 2$ in a circle of radius unity and find the equation of the sphere which has this circle as one of its great circles.

10

2. (a) Let T be the linear operator on R^3 defined by $T(x, y, z) = (2x, 4x - y, 2x + 3y - z)$.

(i) Show that T is invertible.

(ii) Find a formula for T^{-1} . 10

(b) Find the rank of the matrix :

$$A = \begin{pmatrix} 1 & 3 & 1 & -2 & -3 \\ 1 & 4 & 3 & -1 & -4 \\ 2 & 3 & -4 & -7 & -3 \\ 3 & 8 & 1 & -7 & -8 \end{pmatrix} \quad 10$$

(c) Let $A = \begin{pmatrix} 1 & -3 & 3 \\ 3 & -5 & 3 \\ 6 & -6 & 4 \end{pmatrix}$. Is A similar to a diagonal

matrix? If so, find an invertible matrix P such that $P^{-1}AP$ is a diagonal matrix. 10

(d) Find an orthogonal transformation of coordinates to reduce the quadratic form $q(x, y) = 2x^2 + 2xy + 2y^2$ to a canonical form.

10

3. (a) The adiabatic law for the expansion of air is $PV^{1.4} = K$, where K is a constant. If at a given time the volume is observed to be 50 c.c. and the pressure is 30 kg per square centimetre, at what rate is the pressure changing if the volume is decreasing at the rate of 2 c.c. per second? 10

- (b) Determine the asymptotes of the curve $x^3 + x^2y - xy^2 - y^3 + 2xy + 2y^2 - 3x + y = 0$. 10

- (c) Evaluate :

$$\iint_D x \sin(x+y) dx dy,$$

where D is the region bounded by

$$0 \leq x \leq \pi \text{ and } 0 \leq y \leq \frac{\pi}{2}. \quad 10$$

- (d) Evaluate $\iiint (x+y+z+1)^4 dx dy dz$ over the region defined by $x \geq 0$, $y \geq 0$, $z \geq 0$ and $x+y+z \leq 1$. 10

4. (a) Obtain the equations of the planes which pass through the point $(3, 0, 3)$, touch the sphere $x^2 + y^2 + z^2 = 9$ and are parallel to the line $x = 2y = -z$. 10

- (b) The section of a cone whose vertex is P and guiding curve is the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, z = 0$ by the plane $x = 0$ is a rectangular hyperbola.

Show that the locus of P is

$$\frac{x^2}{a^2} + \frac{y^2 + z^2}{b^2} = 1. \quad 10$$

- (c) Prove that the locus of the poles of the tangent planes of the conicoid $ax^2 + by^2 + cz^2 = 1$ with respect to the conicoid $\alpha x^2 + \beta y^2 + \gamma z^2 = 1$ is the conicoid $\frac{\alpha^2 x^2}{a} + \frac{\beta^2 y^2}{b} + \frac{\gamma^2 z^2}{c} = 1$. 10

- (d) Show that the lines drawn from the origin parallel to the normals to the central conicoid $ax^2 + by^2 + cz^2 = 1$ at its points of intersection with the planes $lx + my + nz = p$ generate the cone

$$p^2 \left(\frac{x^2}{a} + \frac{y^2}{b} + \frac{z^2}{c} \right) = \left(\frac{lx}{a} + \frac{my}{b} + \frac{nz}{c} \right)^2. \quad 10$$

Section - B

5. Answer any four :

- (a) Solve :

$$\sec^2 y \frac{dy}{dx} + 2x \tan y = x^3. \quad 10$$

- (b) Find the 2nd order ODE for which e^x and $x^2 e^x$ are solutions. 10

- (c) A uniform rectangular board, whose sides are $2a$ and $2b$, rests in limiting equilibrium in contact with two rough pegs in the same horizontal line at a distance d apart. Show that the inclination θ of the side $2a$ to the horizontal is given by the equation

$$d \cos \lambda [\cos(\lambda + 2\theta)] = a \cos \theta - b \sin \theta$$

where λ is the angle of friction. 10

- (d) A particle rests in equilibrium under the attraction of two centres of force which attract directly as the distance, their intensities being μ and μ' . The particle is slightly displaced towards one of them, show that the time of small oscillation is

$$\frac{2\pi}{\sqrt{(\mu + \mu')}} \quad 10$$

- (e) Verify Green's theorem in the plane for $\oint_C [(xy + y^2)dx + x^2 dy]$ where C is the closed curve of the region bounded by $y = x$ and $y = x^2$. 10

6. (a) Solve :

$$(y^3 - 2yx^2)dx + (2xy^2 - x^3)dy = 0. \quad 10$$

- (b) Solve :

$$\left(\frac{dy}{dx}\right)^2 - 2\frac{dy}{dx} \cos hx + 1 = 0. \quad 8$$

- (c) Solve :

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = x^2 e^{-x}. \quad 10$$

- (d) Show that e^{x^2} is a solution of

$$\frac{d^2y}{dx^2} - 4x\frac{dy}{dx} + (4x^2 - 2)y = 0.$$

Find a second independent solution. 12

7. (a) A solid hemisphere is supported by a string fixed to a point on its rim and to a point on a smooth vertical wall with which the curved surface of the sphere is in contact. If θ and ϕ are the inclinations of the string and the plane base of the hemisphere to the vertical, prove that

$$\tan \phi = \frac{3}{8} + \tan \theta. \quad 10$$

- (b) A particle moves with a central acceleration

$$\mu \left(\gamma + \frac{a^4}{\gamma^3} \right) \text{ being projected from an apse at a}$$

distance a with a velocity $2\sqrt{\mu} a$.

Prove that its path is

$$\gamma^2 (2 + \cos \sqrt{3} \theta) = 3a^2. \quad 10$$

- (c) A shell, lying in a straight smooth horizontal tube, suddenly explodes and breaks into portions of masses m and m' . If d is the distance apart of the masses after a time t , show that the work done by the explosion is

$$\frac{1}{2} \frac{mm'}{m+m'} \frac{d^2}{t^2}. \quad 10$$

- (d) A hollow conical vessel floats in water with its vertex downwards and a certain depth of its axis immersed. When water is poured into it up to the level originally immersed, it sinks till its mouth is on a level with the surface of the water. What portion of axis was originally immersed? 10

8. (a) Show that

$$\bar{A} = (6xy + z^3) \hat{i} + (3x^2 - z) \hat{j} + (3xz^2 - y) \hat{k},$$

is irrotational. Find a scalar function ϕ such that $\bar{A} = \text{grad } \phi$. 10

- (b) Let $\psi(x, y, z)$ be a scalar function. Find $\text{grad } \psi$ and $\nabla^2 \psi$ in spherical coordinates. 8

- (c) Verify Stokes' theorem for

$$\bar{A} = (y - z + 2) \hat{i} + (yz + 4) \hat{j} - xz \hat{k},$$

where S is the surface of the cube $x = 0, y = 0, z = 0, x = 2, y = 2, z = 2$ above the xy -plane.

12

- (d) Show that, if $\bar{r} = x(s) \hat{i} + y(s) \hat{j} + z(s) \hat{k}$ is a

space curve, $\frac{d\bar{r}}{ds} \cdot \frac{d^2\bar{r}}{ds^2} \times \frac{d^3\bar{r}}{ds^3} = \frac{\tau}{\rho^2}$, where τ is

the torsion and ρ the radius of curvature. 10

1.F.5-2009

Sl. No. 8165

B-JGT-J-NBB

MATHEMATICS

Paper—II

Time Allowed : Three Hours

Maximum Marks : 200

INSTRUCTIONS

Candidates should attempt Question Nos. 1 and 5 which are compulsory, and THREE of the remaining questions, selecting at least ONE question from each Section.

The number of marks carried by each question is indicated at the end of the question.

Answers must be written in ENGLISH only.

Assume suitable data, if considered necessary and indicate the same clearly.

Symbols and notations have their usual meanings, unless indicated otherwise.

Section—A

1. Answer any *four* parts from the following :

- (a) Prove that a non-empty subset H of a group G is normal subgroup of $G \Leftrightarrow$ for all $x, y \in H, g \in G, (gx)(gy)^{-1} \in H.$ 10

- (b) Show that the function

$$f(x) = \frac{1}{x}$$

is not uniformly continuous on $]0, 1[.$ 10

- (c) Show that under the transformation

$$w = \frac{z-i}{z+i}$$

real axis in the z -plane is mapped into the circle $|w|=1.$ What portion of the z -plane corresponds to the interior of the circle? 10

- (d) If G is a finite Abelian group, then show that $O(a, b)$ is a divisor of l.c.m. of $O(a), O(b).$ 10

- (e) Evaluate

$$\int_C \frac{2z+1}{z^2+z} dz$$

by Cauchy's integral formula, where C is $|z| = \frac{1}{2}.$ 10

2. (a) Find the dimensions of the largest rectangular parallelepiped that has three faces in the coordinate planes and one vertex in the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1 \quad 14$$

- (b) Determine the analytic function $w = u + iv$, if

$$u = \frac{2\sin 2x}{e^{2y} + e^{-2y} - 2\cos 2x} \quad 13$$

- (c) Find the multiplicative inverse of the element

$$\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$$

of the ring, M , of all matrices of order two over the integers. 13

3. (a) Evaluate

$$\iint xy(x+y) dx dy$$

over the area between $y = x^2$ and $y = x$. 13

- (b) Evaluate by contour integration

$$\int_0^{2\pi} \frac{d\theta}{1 - 2a\sin\theta + a^2}, \quad 0 < a < 1 \quad 13$$

- (c) Write the dual of the following LPP and hence, solve it by graphical method : 14

$$\text{Minimize } Z = 6x_1 + 4x_2$$

constraints

$$2x_1 + x_2 \geq 1$$

$$3x_1 + 4x_2 \geq 1.5$$

$$x_1, x_2 \geq 0$$

4. (a) Show that $d(a) < d(ab)$, where a, b be two non-zero elements of a Euclidean domain R and b is not a unit in R . 13

- (b) Show that a field is an integral domain and a non-zero finite integral domain is a field. 13

- (c) Solve by simplex method, the following LPP : 14

$$\text{Maximize } Z = 5x_1 + 3x_2$$

constraints

$$3x_1 + 5x_2 \leq 15$$

$$5x_1 + 2x_2 \leq 10$$

$$x_1, x_2 \geq 0$$

Section—B

5. Answer any *four* parts from the following :

(a) Find complete and singular integrals of $(p^2 + q^2)y = qz$. 10

(b) Obtain the iterative scheme for finding p th root of a function of single variable using Newton-Raphson method. Hence, find $\sqrt[3]{277234}$ correct to four decimal places. 10

(c) Convert the following binary numbers to the base indicated : 10

(i) $(10111011001.101110)_2$ to octal

(ii) $(10111011001.10111000)_2$ to hexadecimal

(iii) $(0.101)_2$ to decimal

(d) A cannon of mass M , resting on a rough horizontal plane of coefficient of friction μ , is fired with such a charge that the relative velocity of the ball and cannon at the moment when it leaves the cannon is u . Show that the cannon will recoil a distance

$$\left(\frac{mu}{M+m} \right)^2 \frac{1}{2\mu g}$$

along the plane, m being the mass of the ball. 10

- (e) If the velocity of an incompressible fluid at the point (x, y, z) is given by

$$\left(\frac{3xz}{r^5}, \frac{3yz}{r^5}, \frac{3z^2 - r^2}{r^5} \right)$$

where $r^2 = x^2 + y^2 + z^2$, prove that the liquid motion is possible and that the velocity potential is $\cos\theta / r^2$. 10

6. (a) A rod of length l with insulated sides, is initially at a uniform temperature u_0 . Its ends are suddenly cooled to 0°C and are kept at that temperature. Find the temperature distribution in the rod at any time t . 14

- (b) Convert the following to the base indicated against each : 7

(i) $(266.375)_{10}$ to base 8

(ii) $(341.24)_5$ to base 10

(iii) $(43.3125)_{10}$ to base 2

- (c) Draw the circuit diagram for

$$\bar{F} = A\bar{B}C + \bar{C}B$$

using NAND to NAND logic long. 6

- (d) Using Runge-Kutta method, solve $y'' = xy'^2 - y^2$ for $x = 0.2$. Initial conditions are at $x = 0$, $y = 1$ and $y' = 0$. Use four decimal places for computations. 13

7. (a) Prove that the equation of motion of a homogeneous inviscid liquid moving under conservative forces may be written as

$$\frac{\partial \vec{q}}{\partial t} - \vec{q} \times \text{curl } \vec{q} = - \text{grad} \left[\frac{P}{\rho} + \frac{1}{2} q^2 + \vec{\Omega} \right]$$

14

- (b) Find the general solution of

$$\{D^2 - DD' - 2D'^2 + 2D + 2D'\} z \\ = e^{2x+3y} + xy + \sin(2x+y)$$

13

- (c) From the following data

x	:	1	8	27	64
y	:	1	2	3	4

calculate $y(20)$, using Lagrangian interpolation technique. Use four decimal points for computations.

13

8. (a) A homogeneous sphere of radius a , rotating with angular velocity ω about horizontal diameter, is gently placed on a table whose coefficient of friction is μ . Show that there will be slipping at the point of contact for a time $\frac{2\omega a}{7\mu g}$ and that then the sphere will roll with angular velocity $\frac{2\omega}{7}$.

14

- (b) Derive composite $\frac{1}{3}$ rd Simpson's rule.
Hence, evaluate

$$\int_0^{0.6} e^{-x^2} dx$$

by taking seven ordinates. Tabulate the integrand for these ordinates to four decimal places.

13

- (c) Show that for an incompressible steady flow with constant viscosity, the velocity components

$$u(y) = y \frac{U}{h} + \frac{h^2}{2\mu} \left(-\frac{dp}{dx} \right) \frac{y}{h} \left(1 - \frac{y}{h} \right),$$

$$v = 0, w = 0$$

satisfy the equations of motion, when the body force is neglected. $h, U, \frac{dp}{dx}$ are constants and $p = p(x)$.

13
